

Undernutrition & childhood morbidities among tribal preschool children

V. G. Rao, Rajeev Yadav, C.K. Dolla, Surendra Kumar, M.K. Bhoneley & Mahendra Ukey

Regional Medical Research Centre for Tribals (ICMR), Jabalpur, India

Received July 22, 2004

Background & objectives: Undernutrition and various morbidities go hand in hand, particularly in children. Nutritional status is a sensitive indicator of community health and nutrition. The present study is an attempt to assess the nutritional status of pre-school children of Gond tribal community in Madhya Pradesh.

Methods: The study was a community-based, cross-sectional survey carried out in tribal pre-school children. Anthropometric measurements were taken. Various indices of nutritional status were expressed in standard deviation units (z scores) from the reference median. The children were examined for nutritional deficiencies and other morbidities. The haemoglobin concentration was measured and the children were classified into various grades of nutritional anaemias. Data on socio-cultural and hygienic practices were also collected.

Results: More than 60 per cent children were underweight. Micronutrient deficiency disorders such as anaemia and vitamin A deficiency were common among them. Unhygienic personal habits and adverse cultural practices relating to child rearing, breast-feeding and weaning were also prevalent among them.

Interpretation & conclusion: The findings of the present study revealed the widespread prevalence of undernutrition among pre-school tribal children and highlight a need for an integrated approach towards improving the child health as well as nutritional status in this area.

Key words Anaemia - pre-school children - tribal - undernutrition - unhygienic practices

Pre-school children constitute the most vulnerable segment of any community. Their nutritional status is a sensitive indicator of community health and nutrition¹. Undernutrition among them is one of the greatest public health problems in developing countries. About 128 million (70%) of the world's 182 million stunted children aged under five years live in Asia².

Nutritional status plays a vital role in deciding the health status particularly in children. Nutritional deficiencies give rise to various morbidities, which in turn, may lead to increased mortality. Undernutrition is a known factor closely associated with child mortality rates^{3,4}. An analysis of six longitudinal studies by World Health Organization (WHO) revealed a strong association between severity of weight for

age deficits and mortality rates: 54 per cent deaths of under five children in developing countries were accompanied by low weight for age⁵. Attempts to reduce child mortality in developing countries through selective primary health care have focused primarily on the prevention and control of specific infectious diseases, with less effort being directed to improving children's underlying nutritional status⁶.

The subject gains more importance in tribal children, due to certain adverse realities like insufficient food intake, frequent infections, lack of access to health services, illiteracy, unhygienic personal habits, adverse cultural practices, *etc.* In this study we made an attempt towards assessing the nutritional status, nutritional deficiency signs and other morbidities among pre-school children of *Gond* tribal community of Madhya Pradesh, India.

Material & Methods

The study was carried out in Kundam block of Jabalpur district, Madhya Pradesh (MP), during April 2000 and May 2001. It was a community based, cross-sectional survey among *Gond* tribe of the area. Majority of the villages in the block are inhabited by the *Gond* tribal population. In order to assess nutritional status of pre-school children, 27 of the 197 tribal villages were selected by the probability proportionate to size (PPS) sampling.

A house-to-house survey was conducted in study villages to examine tribal pre-school children. The sample size was estimated to be 817 at 95 per cent confidence interval (CI) with 5 per cent precision. However, considering anticipated non response in the field situation, oversampling of 25 per cent was made to achieve the estimated sample size. Thus, a total of 1022 children were covered from 27 villages. Anthropometric measurements in the form of height and weight were taken using standard techniques⁵. Height of children under two years of age was measured in supine position using infantometer (Seca, Germany) and older children were measured using infantometer. The body weight of each child was measured using lever balance. The ages of the children were obtained from the birth records available with the village panchayat and were cross-checked with the parents by using local events calendar to ensure accuracy of the records. The

heights and weights of each child were compared with National Center for Health Statistics (NCHS) reference data⁷ for age and sex of children. The indices of nutritional status (weight for age, height for age and weight for height) were expressed in standard deviation units (z scores) from the reference median as recommended by the WHO⁸. Children with more than two standard deviations below the reference median on any of the indices were considered to be undernourished and the children who fell more than three standard deviations below the reference median were considered to be severely undernourished. All the children were examined clinically by a medical epidemiologist for clinical signs of nutritional deficiencies and other morbidities. The haemoglobin concentration was measured using Sahli's method⁹. Children were classified into different grades of nutritional anaemia as per the WHO criteria¹⁰.

Data on socio-cultural and hygienic practices were collected from the households during the survey using pre-tested, structured proforma. Data were analyzed using t test and χ^2 test with the help of Statistical Package for Social Sciences (SPSS) software.

Results

A total of 1022 pre-school children (M 527, F 495) were examined. Boys were slightly taller and heavier than girls but the difference was not significant (Table I). Both the boys and girls were shorter and lighter compared to the NCHS reference data for their ages and sex.

Table II shows distribution of pre-school children according to Standard Deviation (SD) classification. High prevalence of undernutrition (below 2SD) in terms of underweight (61.6%), stunting (51.6%) and wasting (32.9%) was observed among them. The study revealed severe degree (below -3SD) of underweight, stunting and wasting in 27.8, 30.3 and 6.5 per cent children, respectively. Prevalence of undernutrition in terms of underweight, stunting and wasting was found similar in both the sexes (Table III).

Anaemia prevalence rate of 86.7 per cent was observed among the children. Moderate and severe anaemia was observed in 71.1 per cent children. Prevalence and the distribution of various grades of anaemia were found similar in boys as well as in girls (Table IV).

Table I. Mean anthropometric measurements of pre-school children

Age (yr)	Sex	N	Weight (kg)	Height (cm)
0-1	M	75	6.2 ± 1.75	63.6 ± 4.86
	F	69	5.8 ± 1.68	62.8 ± 5.02
1+	M	85	8.2 ± 1.64	72.3 ± 5.48
	F	82	7.9 ± 1.53	70.9 ± 4.95
2+	M	94	9.8 ± 1.81	82.0 ± 5.32
	F	88	9.5 ± 1.60	80.1 ± 5.10
3+	M	89	11.2 ± 1.72	86.2 ± 6.28
	F	86	11.0 ± 1.56	85.6 ± 5.51
4+	M	96	12.8 ± 1.98	95.4 ± 5.56
	F	89	12.4 ± 1.91	94.3 ± 6.82
5+	M	88	13.8 ± 2.01	101.2 ± 6.19
	F	81	13.6 ± 2.22	99.7 ± 6.77

Values are given as mean ± SD

Prevalence of clinical protein energy malnutrition (PEM) in the form of marasmus was found in 6 (0.6%) children, while Kwashiorkor was not recorded. Emaciation was seen in 35 (3.4%) children. Clinical signs of anaemia were present in 530 (51.9%) children. Vitamin A deficiency in the form of Bitot’s spots was recorded in 16 (1.6%) children. A total of 127 (12.4%) children were found suffering from upper respiratory tract infection. Vitamin B complex deficiency in the form of angular stomatitis and cheilosis was seen in 29 (2.8%) children.

Data on socio-cultural and hygienic practices revealed that open air defecation is the rule in these villages. Children often defecate in the open space around the houses. Most of them either do not wash their hands or use plain water for hand washing before eating and after defecation. Only 93 (9.1%) children were put on breast feeding within 2 h of birth; 380 (37.2%) children were put to breast by second day or

thereafter. Majority of the children (72.5%) did not receive prelacteal feeds. Exclusive breast feeding up to 4 months was not done in 23.4 per cent children. Supplementary feeding was, however, started by 6 months in majority of the children (84.5%). Colostrum was not given to 86.4 per cent children.

Discussion

Undernutrition was found to be widely prevalent among pre-school children of the *Gond* tribe of Jabalpur district. More than 60 per cent children were underweight and more than 25 per cent had severe undernutrition. High prevalence of chronic undernutrition was also observed in other studies¹¹⁻¹⁴. Higher prevalence of undernutrition was also observed among them as compared to the prevalence among children in general^{15,16}. The findings of the National Family Health Survey (NFHS) also reveals the highest proportion of underweight children in Madhya Pradesh as compared to other states in India, and according to other two indicators also, higher prevalence of undernutrition was observed in the state of Madhya Pradesh¹⁷.

Contrary to popular belief, we found that colostrum was not given to majority of the children and exclusive breast-feeding up to 4 months was not done in about one fourth of the children. More than one third of the children were put to breast by second day of birth or thereafter. Giving prelacteal feed was lower in tribal children than in other urban studies^{18,19}.

Undernutrition is a well known contributory factor to high mortality in children due to infectious diseases. High infant mortality and under five mortality rates among scheduled tribes of Madhya Pradesh have been observed¹⁷. Though the causes of high rate of childhood mortality were not studied, high prevalence of respiratory tract infections and

Table II. Distribution of pre-school children according to SD classification

	Pre-school children				
	<-3SD	-3SD to <-2SD	-2SD to <-1SD	-1SD to <median	≥median
Weight/age	283 (27.7)	347 (33.9)	240 (23.5)	93 (9.1)	59 (5.8)
Height/age	308 (30.1)	220 (21.5)	200 (19.6)	155 (15.2)	139 (13.6)
Weight/height	66 (6.5)	270 (26.4)	419 (41.0)	198 (19.4)	69 (6.7)

Figures in parentheses indicate percentages

Table III. Nutritional status of pre-school children

Children	Underweight (%)	Stunting (%)	Wasting (%)
Boys (n=527)	332 (62.9)	287 (54.4)	159 (30.1)
Girls (n=495)	298 (60.3)	241 (48.8)	177 (35.8)
Total (n=1022)	630 (61.6)	528 (51.6)	336 (32.9)

t(a), (b) and (C): $P > 0.05$

Table IV. Prevalence of anaemia among pre-school children

Anaemia status	Boys (n=509)	Girls (n=476)	Total (n=985)
Mild	86 (16.9)	68 (14.3)	154 (15.6)
Moderate	313 (61.5)	315 (66.2)	628 (63.8)
Severe	39 (7.7)	33 (6.9)	72 (7.3)
Total	438 (86.1)	416 (87.4)	854 (86.7)

Figures in parentheses indicate percentages

$\chi^2=2.04$ at 2df $P > 0.08$

micronutrient deficiency disorders such as anaemia and vitamin A deficiency might be playing an important role. A study among preschool children of this tribe also revealed high prevalence of intestinal parasitic infections (48%) among them²⁰. Poor environmental sanitation and unhygienic personal habits appear to predispose them to the risk of infections. In addition to the infections, adverse cultural practices relating to child rearing, breast-feeding and weaning were some other contributory factors for malnutrition and must be playing an important role in this community.

The present study revealed the widespread prevalence of undernutrition among pre-school children of this tribal community. Analysis of pooled data obtained from several studies has shown that children with mild to moderate degrees of malnutrition have a relative risk of 2.2 for death under five years of age and children with severe malnutrition have a relative risk of 6.8²¹. Also, the population attributable risk for death due to potentiating effect of malnutrition on infectious diseases varies with the percentage of children, with weight for age less than 80 per cent of the reference point, following a quadratic relation³ and 21 per cent of all deaths occurring among under-fives is attributable to malnutrition²⁰.

Though undernutrition *per se* cannot be the direct cause of high rate of childhood mortality it appears to be a major contributory factor for high infant and child mortality rates. The problem of undernutrition amongst pre-school tribal children needs to be addressed through comprehensive preventive, promotive and curative measures. The community needs to be educated about environmental sanitation and personal hygienic practices, and also proper child rearing, breast-feeding and weaning practices. Appropriate nutritional programmes should be designed to meet the requirements. A comprehensive child survival programme with supplementary feeding, growth and development monitoring and early, prompt treatment during illness needs to be devised and implemented ensuring community participation.

Acknowledgment

The authors thank Prof. A.P. Dash, Director, Regional Medical Research Centre for Tribals (ICMR), Jabalpur for his constant support and guidance. The technical support and cooperation given by Mr Sujeet Das, Mr J.P. Mishra, Mr R.K. Minocha, Mr J.P. Misra, Mr R.K. Minocha, Mr Lalit Sahare, Mr Ajesh Dubey and Mr Rajendra Kusare is gratefully acknowledged.

References

1. Sachdev HPS. Assessing child malnutrition - some basic issues. *Bull Nutr Foundations India* 1995; 16 : 1-5.
2. Allen LH, Gillespie SR. What works? A review of the efficacy and effectiveness of nutrition interventions. United Nations. Administrative Committee on Coordination/Sub-Committee on Nutrition, Geneva. Asian Development Bank, Manila. ACC/SCN Nutrition Policy Paper No.19. *ADB Nutrition and Development series Geneva* 2001 No.5, p. 24-5.
3. Pellitier DL, Frongillo EA Jr, Schroeder DG, Habicht JP. The effects of malnutrition on child mortality in developing countries. *Bull World Health Organ* 1995; 73 : 443-8.
4. Scrimshaw NS, Taylor CE, Gordon JE. Interactions of nutrition and infection. WHO Monograph Series 57, Geneva, World Health Organization; 1968 p. 3-329
5. (No authors listed). Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organ Tech Rep Ser* 1995; 854: 1-452.

6. Walsh JA, Warren KS. Selective primary health care - an interim strategy for disease control in developing countries. *N Engl J Med* 1979; 301 : 967-74.
7. World Health Organization. *Measuring change in nutritional status*. Geneva: World Health Organization; 1983 p. 64-100.
8. (No authors listed). Use and interpretation of anthropometric indicators of nutritional status. WHO Working Group. *Bull World Health Organ* 1986; 64 : 929-41.
9. Godkar P. *Text book of medical laboratory technology*. Mumbai: Bhalani Publishing House; 2001 p. 439.
10. (No authors listed). Nutritional anaemias: Report of a WHO Group of Experts. *WHO Tech Rep Ser* 1972; 503: 1-29.
11. Iqbal Hossain M, Yasmin R, Kabir I. Nutritional and immunisation status, weaning practices and socioeconomic conditions of under five children in three villages of Bangladesh. *Indian J Public Health* 1999; 43 : 37-41.
12. Mahapatra A, Geddam JJB, Marai N, Murmu B, Mallick G, Bulliyya G, *et al.* Nutritional status of preschool children in the drought affected Kalahandi district of Orissa. *Indian J Med Res* 2000; 111 : 90-4.
13. Sood AK, Kapil U. Anthropometry in detection of protein energy malnutrition. *Indian Pediatr* 1984; 21 : 635-9.
14. National Nutrition Monitoring Bureau. Diet and nutritional status of tribal population - Report on first repeat survey (1998-99). Hyderabad: National Institute of Nutrition; 2000 p. 14.
15. International Institute for Population Sciences (IIPS) and ORC Macro. 2000. *National Family Health Survey (NFHS-2), 1998-99: India*. Mumbai, IIPS.
16. UNICEF: *The state of the World's children*. Girls, education and development. New York: UNICEF 2004 p. 107.
17. International Institute for Population Sciences (IIPS) and ORC Macro. 2001. *National Family Health Survey (NFHS-2) , India, 1998-99: Madhya Pradesh*.
18. Bahl L, Kaushal RK. Infant rearing practices & beliefs in rural inhabitants of Himachal Pradesh. *Indian Pediatr* 1987; 24 : 903-6.
19. Kar M, De R. Breast feeding practices. Impressions from an urban community. *Indian J Public Health* 1991; 35 : 93-6.
20. Rao VG, Yadav R, Bhondeley MK, Das S, Agrawal MC, Tiwary RS. Worm infestation and anaemia: a public health problem among tribal pre-school children of Madhya Pradesh. *J Commun Dis* 2002; 34 : 100-5.
21. Schroeder DG, Brown KH. Nutritional status as a predictor of child survival - Summarizing the association and quantifying its global impact. *Bull World Health Organ* 1994; 72 : 569-79.

Reprint requests: Dr V.G. Rao, Deputy Director, Regional Medical Research Centre for Tribals (ICMR)
Nagpur Road, P.O. Gama, Jabalpur 482003, India
e-mail: drvgrao@rediffmail.com

Malnutrition among children is rampant among the developing countries. About half of all children deaths are associated with malnutrition, of which three quarters are linked to mild and moderate forms (Ruwali, 2011) [12]. Widespread poverty, illiteracy, malnutrition, poor maternal and child health services and ineffective coverage of national health and nutritional services have been traced out in several studies as possible contributing factors to dismal health conditions prevailing among the tribal population in India. Undernutrition and various morbidities go hand in hand, particularly in children. Nutritional status is a sensitive indicator among pre-school tribal children and highlight a need for an integrated approach towards improving the child health as. Childhood undernutrition is a global health challenge. Undernutrition in early life is associated with a number of adverse outcomes, including persistent stunting, immune dysfunction, and neurocognitive deficits. Current approaches to treatment have only modest effects in correcting these long-term sequelae, suggesting that certain features of host biology are not being adequately repaired. Compared with healthy children, microbiota from undernourished children transmit impaired growth phenotypes to recipient gnotobiotic mice fed diets representative of those of the human donors; moreover, some of the transplanted age-discriminatory strains are growth-discriminatory.

@article{Rao2005UndernutritionC, title={Undernutrition & childhood morbidities among tribal preschool children.}, author={Vikas G Rao and Rajeev Ranjan Yadav and Chandra Kumar Dolla and Surendra Kumar and Manoj Kumar Bhondeley and Mahendra Ukey}, journal={The Indian journal of medical research}, year={2005}, volume={122 1}, pages={43-7 } }. Vikas G Rao, Rajeev Ranjan Yadav, +3 authors Mahendra Ukey. BACKGROUND & OBJECTIVE Undernutrition and various morbidities go hand in hand, particularly in children. Nutritional status is a sensitive indicator of community health and nutrition. The